

- Plan:
1. Implicit differentiation
 2. The second derivative and curvature
 3. Convex optimization

1. Implicit differentiation

Ex A curve is implicitly defined by
the equation $y^2 - x^3 = 1$

- Express y' by x and y using implicit differentiation
- Find all solutions for y when $x = 2$
- compute y' for these points
- Find the function expression(s) for the tangent line(s) at $x = 2$.

Solution

a) $(y^2)'_x - (x^3)'_x = (1)'_x$

Chain rule: $u = y$ and $g(u) = u^2$

$$u'_x = y'_x \quad g'(u) = 2u$$

$$\text{so } (y^2)'_x = 2u \cdot u'_x = 2yy'$$

$$2yy' - 3x^2 = 0 \quad \begin{matrix} \text{- solve for } y' \\ 2y \cdot y' = 3x^2 \end{matrix}$$

$$y' = \frac{3x^2}{2y}$$

b) $x=2$, solve $y^2 - 2^3 = 1$ for y

$$y^2 = 1 + 8 = 9$$
$$\underline{\underline{y = \pm 3}}$$

c) $(2, 3)$: $y' = \frac{3 \cdot 2^2}{2 \cdot 3} = \underline{\underline{2}}$

$(2, -3)$: $y' = \frac{3 \cdot 2^2}{2 \cdot (-3)} = \underline{\underline{-2}}$

d) Tangent line through $(2, 3)$:


$$h(x) - 3 = 2(x - 2)$$

so $\underline{\underline{h(x) = 2x - 1}}$

Tangent line through $(2, -3)$


$$g(x) - (-3) = -2 \cdot (x - 2)$$

so $\underline{\underline{g(x) = -2x + 1}}$




2. The second order derivative and curvature

bending up




3 graphs of
convex functions

bending down



3 graphs of
concave functions



Definition

- $f(x)$ is convex in the interval $[a, b]$ if $f''(x) \geq 0$ for all x in (a, b) .
- $f(x)$ is concave in the interval $[a, b]$ if $f''(x) \leq 0$ for all x in (a, b) .
- A number c is an inflection point for $f(x)$ if $f''(x)$ changes sign at $x=c$.

Start: 11.10

③


Note If $f(x)$ is convex, then $f'(x)$ is an increasing function.

If $f(x)$ is concave then $f'(x)$ is a decreasing function.


Second derivative test (sec. 8.5)

Suppose $x = a$ is a stationary point for $f(x)$.

If $f''(a) > 0$ then $x = a$ is a (local) minimum point



If $f''(a) < 0$ then $x = a$ is a (local) maximum point.



Ex $f(x) = x^3 - 3x^2 + 5$. Find local max/min. points by using the second derivative test.

Solution

$f'(x) = 3x^2 - 6x$ and find the stationary points as the solutions to the eq.

$$3x^2 - 6x = 0$$

(4)

$3x$ is a common factor

$$3x(x - 2) = 0$$

either $x = 0$ or $x = 2$

Calculate $f''(x) = [f'(x)]' = (3x^2 - 6x)'$
 $= 6x - 6$

$$f''(0) = 6 \cdot 0 - 6 = -6 < 0$$

so $x = 0$ is a (local) maximum point

$$f''(2) = 6 \cdot 2 - 6 = 6 > 0$$

so $x = 2$ is a (local) minimum point.

3. Convex optimization

Fact If $f(x)$ is convex everywhere in its domain, then any stationary point is a global minimum point.

And if $f(x)$ is concave everywhere in its domain (an interval), then any stationary point is a global max. point.

Ex $f(x) = x^4 + 5x^2 + 3$, $D_f = \leftarrow, \rightarrow\right\rangle$

- a) Find the stationary points $= \langle \infty, \infty \rangle = \mathbb{R}$.
- b) Determine if they are global max. or min. pts.
- c) \longrightarrow the extremal values.

Solution

a) calculate $f'(x) = 4x^3 + 10x$

Stationary points are solutions to the eq $4x^3 + 10x = 0$ (x is a common factor)

$$x(4x^2 + 10) = 0$$

so $\underline{x = 0}$ is the only solution.

b) calculate $f''(x) = 12x^2 + 10$

which is greater or equal to 10 for all x . So $x = 0$ is a global min. point.

c) $f(0) = 0^4 + 5 \cdot 0^3 + 3 = \underline{3}$ is the global minimal value.