

Solutions:

Key problems Lecture 11

1. a) $3t^2 - 2t + 2y \cdot y' = 0$

Exact? $p = 3t^2 - 2t = h'_t$
 $q = 2y = h'_y$

II

I: $h = \int 3t^2 - 2t \, dt = t^3 - t^2 + g(y)$

II: $h'_y = 0 + g'(y) = 2y \Rightarrow g(y) = y^2 + C = y^2 \leftarrow \begin{matrix} \text{may} \\ \text{choose} \\ C=0 \end{matrix}$

II

$h = t^3 - t^2 + y^2$ solution I-II, eqn. is exact

Solution: $t^3 - t^2 + y^2 = C \leftarrow h(t,y) = C$

$$y^2 = C + t^2 - t^3$$

$$y = \pm \sqrt{C + t^2 - t^3}$$

b) $2y - 3t^2 + 2(y+t)y' = 0$

$p = 2y - 3t^2 = h'_t \Rightarrow h = 2yt - t^3 + g(y)$

$q = 2y + 2t = h'_y \quad h'_y = 2t - 0 + g'(y) = 2y + 2t$

$$g'(y) = 2y$$

$$g(y) = y^2 + C = y^2$$

$h = 2yt - t^3 + y^2$ exact

$$2yt - t^3 + y^2 = C$$

$$y^2 + 2t \cdot y + (-t^3 - C) = 0$$

$$y = \frac{-2t \pm \sqrt{4t^2 - 4 \cdot (-t^3 - C)}}{2}$$

$$= -t \pm \sqrt{t^2 + t^3 + C}$$

$$c) \frac{y(1-2\ln t)}{t^3} + \frac{\ln t}{t^2} y' = 0$$

$$p = y \frac{(1-2\ln t)}{t^3} = h'_t$$

$$q = \frac{\ln t}{t^2} = h'_y$$

Start with second eqn:

$$\Rightarrow h = \frac{\ln t}{t^2} \cdot y + g(t)$$

$$h'_t = \frac{\frac{1}{t} \cdot t^2 - \ln t \cdot 2t}{t^4} y + g'(t)$$

$$= \frac{t - 2t \ln t}{t^4} y + g'(t)$$

$$p = y \frac{(1-2\ln t)}{t^3}$$

$$\xleftarrow[\substack{\text{should} \\ \text{equal} \\ p}]{=} \frac{1 - 2\ln t}{t^3} y + g'(t)$$

ok if $g'(t) = 0$


choose $g(t) = 0$

$$h = \frac{\ln t}{t^2} \cdot y = c$$

$$y = \underline{\underline{\frac{ct^2}{\ln t}}}$$


2. a) $y' = 6 - 2y$

Eq. state: $6 - 2y = 0$
 $y = 3 \Rightarrow y_e = 3$



phase diagram

$y_e = 3$ is globally asymptotically stable




b) $y' = y^2 - 4$

Eq. state: $y^2 - 4 = 0$
 $y = \pm 2$


Eq. states:

$y_e = -2$ and $y_e = 2$



$y_e = 2$ is unstable

$y_e = -2$ is stable, but not globally asymptotically stable



Alternative method: Stability thm.

$F(y) = y^2 - 4$

$F'(y) = 2y$

$F'(2) = 4 > 0$

$F'(-2) = -4 < 0$

$\Rightarrow y_e = 2$ unstable

$y_e = -2$ stable

note: cannot tell if $y_e = -2$ is globally asymptotically stable from $F'(-2)$.


$$c) y' = sy(1 - y/10)$$

Eg. state:

$$sy(1 - y/10) = 0$$

$$y=0 \text{ or } 1 - y/10 = 0$$

$$y=10$$




phase diagram

$y_c=0$ unstable

$y_c=10$ stable, but not globally asymptotically stable

Eg. states:

$$\underline{y_c=0} \text{ and } \underline{y_c=10}$$



3. a) $y'' + 6y' - 16y = 16t - 22$

$$y = y_h + y_p = \underline{\underline{C_1 e^{2t} + C_2 e^{-8t} - t + 1}}$$

y_h : $y'' + 6y' - 16y = 0$

$$r^2 + 6r - 16 = 0$$

$$r = 2, r = -8 \Rightarrow y_h = \underline{\underline{C_1 e^{2t} + C_2 e^{-8t}}}$$

y_p : $y'' + 6y' - 16y = 16t - 22$ $\leftarrow f(t) = 16t - 22$
 $f'(t) = 16$
 $f''(t) = 0$

$$0 + 6 \cdot A - 16(At + B) = 16t - 22$$

$$(-16A)t + (6A - 16B) = 16t - 22$$

$\uparrow$
coeff. of t

$\uparrow$
const.

$$\left\{ \begin{array}{l} \text{Guess:} \\ y = At + B \\ y' = A \\ y'' = 0 \end{array} \right.$$

Comparison of coeff's:

$$-16A = 16 \Rightarrow A = \underline{\underline{-1}}$$

$$6A - 16B = -22$$

$$-6 - 16B = -22$$

$$-16B = -16$$

$$\underline{\underline{B = 1}}$$

$$y_p = At + B = \underline{\underline{-t + 1}}$$

$$b) \quad y'' + 6y' + 9y = 4e^{-t}$$

$$y = y_h + y_p = \underline{\underline{C_1 e^{-3t} + C_2 t e^{-3t} + e^{-t}}}$$

$$y_h: \quad y'' + 6y' + 9y = 0$$

$$r^2 + 6r + 9 = 0$$

$$r = \frac{-6 \pm \sqrt{36 - 4 \cdot 9}}{2}$$

$$= -3 \pm 0$$

$$r_1 = r_2 = \underline{-3} \text{ (double root)} \Rightarrow y_h = \underline{\underline{C_1 e^{-3t} + C_2 t e^{-3t}}}$$

$$y_p: \quad y'' + 6y' + 9y = 4e^{-t}$$

$$f = 4e^{-t}$$

$$f' = -4e^{-t}$$

$$f'' = 4e^{-t}$$

$$y = A e^{-t} \leftarrow \underline{\text{guess}}$$

$$y' = -A e^{-t}$$

$$y'' = A e^{-t}$$

$$(A e^{-t}) + 6(-A e^{-t})$$

$$+ 9(A e^{-t}) = 4e^{-t}$$

$$(A - 6A + 9A) e^{-t} = 4e^{-t}$$

$$(4A) e^{-t} = 4e^{-t}$$

Comparison of coeff's:

$$4A = 4$$

$$A = 1$$

$$y_p = A e^{-t} = \underline{\underline{e^{-t}}}$$

$$c) \quad y'' - 3y' + 2y = 3e^{2t}$$

$$y = y_h + y_p = \underline{C_1 e^t + C_2 e^{2t} + 3te^{2t}}$$

$$y_h: \quad y'' - 3y' + 2y = 0$$

$$r^2 - 3r + 2 = 0$$

$$r=1, r=2$$

$$\Rightarrow y_h = \underline{C_1 e^t + C_2 e^{2t}}$$

$$y_p: \quad y'' - 3y' + 2y = 3e^{2t}$$

$$f = 3e^{2t}$$

$$f' = 6e^{2t}$$

$$f'' = 12e^{2t}$$

$$y = Ae^{2t} \leftarrow \text{guess}$$

$$y' = 2Ae^{2t}$$

$$y'' = 4Ae^{2t}$$

$$(4Ae^{2t}) - 3(2Ae^{2t}) + 2(Ae^{2t}) = 3e^{2t}$$

$$(4A - 6A + 2A)e^{2t} = 3e^{2t}$$

$$0Ae^{2t} = 3e^{2t} \leftarrow \text{no solution}$$

$$(4A + 4At)e^{2t} - 3(A + 2At)e^{2t} + 2 \cdot (At)e^{2t} = 3e^{2t}$$

$$(4A - 3A + 4At - 6At + 2At)e^{2t} = 3e^{2t}$$

$$Ae^{2t} = 3e^{2t}$$

$$\underline{A=3}$$

$$y_p = \underline{Ate^{2t} = 3te^{2t}}$$

try

$$y = t \cdot Ae^{2t} = \underline{Ate^{2t}}$$

$$y' = Ae^{2t} + At \cdot e^{2t} \cdot 2 = \underline{(A + 2At)e^{2t}}$$

$$y'' = 2Ae^{2t} + (A + 2At)e^{2t} \cdot 2 = \underline{(4A + 4At)e^{2t}}$$

$$d) y'' - y = t^2$$

$$y = y_h + y_p = \underline{C_1 e^t + C_2 e^{-t} - t^2 - 2}$$

$$\underline{y_h}: y'' - y = 0$$

$$r^2 - 1 = 0$$

$$r = \pm 1$$

$$\Rightarrow y_h = \underline{C_1 e^t + C_2 e^{-t}}$$

$$\underline{y_p}: y'' - y = \boxed{t^2}$$

$$f = t^2$$

$$f' = 2t$$

$$f'' = 2$$

$$\underline{\text{Guess:}} \quad y = At^2 + Bt + C$$

$$y' = 2At + B$$

$$y'' = 2A$$



$$2A - (At^2 + Bt + C) = t^2$$

$$(-A)t^2 + (-B)t + (2A - C) = t^2$$

$$\underline{\text{Comparison of coeff's:}} \quad (t^2 = 1 \cdot t^2 + 0 \cdot t + 0 \cdot 1)$$

$$-A = 1 \quad \underline{A = -1}$$

$$-B = 0 \quad \underline{B = 0}$$

$$2A - C = 0 \quad \underline{C = 2A = -2}$$

$$y_p = At^2 + Bt + C = \underline{-t^2 - 2}$$